

# Trace the Denmark Strait overflow water in an eddy-resolving Atlantic simulation: some preliminary results

Xiaobiao Xu (COAPS/FSU),  
Alan Wallcraft (NRL/SSC),  
Eric Chassignet (COAPS/FSU)


Thanks: Peter Rhines (UW) and William Schmitz

May 21-23, Layered ocean modeling workshop, Ann Arbor, MI

# Motivation


- Passive tracer, when tagged to a particular water mass, can be used to “effectively” track the movement/circulation of the water mass and its interaction with the ambient water.
- When the Denmark Strait overflow water (DSOW) flow into the Irminger and Labrador Seas, the observed vertical structure of horizontal velocity shows a strong bottom intensification, typically not in general circulation models. We want to use tracer to quantify the associated mixing and water mass transformation.
- Tracers (CFC, Oxygen,  $I^{129}$ , etc) have been widely used in observations associated with the DSOW, e.g., Smith et al., 2005. Can models simulate those tracer signature?

# 3-layer circulation in the SPNA




upper layer: Red and yellow (“modified” North Atlantic Water); middle layer: green (Labrador Sea Water); deep layer: blue (Nordic Seas Overflow Water)

# Overflow structure in HYCOM vs. obs.




(a) Angmagssalik



(b) Southeast of Cape Farewell


# Comparison near 53N

Average mooring velocities and transports (July 97 July 99)



Moored observations from Fischer et al., 2004, 2010. HYCOM results and comparison to observations see Xu et al., 2013

# Comparison near 43N




Moored observations from Clarke et al., 1998; Meinen 2000; Schott et al., 2004, 2006. Model results and comparison to observations see Xu et al., 2013

# Tracer in the Atlantic Simulation


- HYbrid Coordinate Ocean Model (HYCOM, Bleck, 2002; Chassignet et al., 2003; <http://www.hycom.org>); see Xu et al., 2010, 2012, 2013 for documentation of the Atlantic model results
- Model domain extends meridionally from 28° S to 80° N, as a subset of the global prediction system developed at NRL/SSC
- 0.08° horizontal resolution (5-6 km in the SPNA); 32  $\sigma_2$  layers in vertical, with thermo-baric effect parameterized
- Initialization with zero velocity and January potential temperature and salinity from monthly ocean climatology GDEM (Carnes, 2009)
- Climatological forcing (E026): 25-year spin-up using forcing based on a monthly climatology (ERA40, Uppala et al., 2005) + submonthly wind anomalies from Navy Operational Global Atmospheric Prediction System (NOGAPS) for year 2003;
- Tracers were injected on Jan 1 of model year 16 (at the Denmark Strait sill below model layer 21 and in the Florida Strait in the top 15 layers). The simulation is integrated for 10 years.

# 10 years after (Florida Strait)



Integrated tracer content below model layers at the end of the simulation;

# 10 years after (Denmark Strait)









Integrated tracer content below model layer 20 at the end of the simulation;






# Vertical distribution (1 year mean)

Passive-tracer 67.88n - 65.45n mean: 23.955- 25.045 [03.2H]

Passive-tracer 53.67n - 60.51n mean: 23.955- 25.045 [03.2H]




# Compared to the observed $I^{129}$




# In the Labrador Sea

I129 from Smith et al., 2005




Both the observed I<sup>129</sup> and numerical tracer spread out the lowest part of *entire* Labrador Sea.




# Vertical structure of tracer flux


## Denark Strait




## Angmagssalik




## Cape Farewell




## Labrador Sea




## Grand Banks





## 50W




# Total tracer fluxes in overflow water




# Full water column budget



**Net flux through the boundaries**







**Inventory change in the box**



# Summary

- A numerical tracer experiment is conducted in an eddy-resolving Atlantic simulation.
- The DSOW tracer is seen descend from the Denmark Sill to the observed DSOW depth range in the Irminger and Labrador Sea, with a vertical distribution similar to the observed  $I^{129}$ .
- Preliminary analysis suggest that immediately after the DSOW spreading into the Irminger Sea, some tracer have been *lost* to water above the overflow.
- The full water column budget of the tracer flux/content is *approximately* conserved.
- **\*\*currently on going \*\*** simplify the experiment by releasing tracer only in the DSOW, and save the model fluxes to test the conservation of tracer content.

# If I saved the model volume fluxes ...



# Thank you

